




1

CMTS DATA Support


 Six-month MOUD summaries- data audit
 Please faithfully enter no-shows for you and your prescribers going forward
 Please monitor that you've entering the MOUD medication accurately on the prescriber encounter


WMedicine



3









Buprenorphine



- Partial agonist with slow onset
- Available as tablet or film, generally offered in a duo product with Naloxone (Narcan) for safety
- Effect on the brain
- Ceiling effect on respiratory depression
- High affinity for μ -opioid receptor
- Slowly dissociates from μ-opioid receptors

UW Medicine

Buprenorphine Side Effects



- Side effects may be less intense than full agonists
- Minimal potential for respiratory depression Safer than
- Patients may experience oral issues at site of ingestion (dissolves in the mouth)
- Other typical opioid side effects:
- Constipation, Vomiting
 Insomnia/sleepiness

- Patients' principal concern may be withdrawal effects in transition from street drugs to buprenorphine
 - Importance of comfort meds

UW Medicine

9

10

Main Buprenorphine Products





- Suboxone (Zubsolv, Bunavail) "bupe, subs"
- Buprenorphine with naloxone (Narcan)
- Most common medication formulation
- **Sublocade** (Extended-Release Buprenorphine)
 - Monthly injection
 - Indicated for patients who have been stable on buprenorphine for at least 7 days
- Subutex
 - Buprenorphine mono product- higher risk for abuse/diversion


LW Medicine

CHAMP

Current Issues in Street Opioids

LW Medicine

11



Fentanyl- a extremely powerful drug



- Fentanyl was involved in 64% of >100,000 estimated U.S. drug overdose deaths during May 2020–April 2021
- Injection drug use was reported in 25% of deaths, and non-injection routes of drug use in **27%** of deaths
- 40% of Fentanyl-involved deaths also involved stimulants

O'Donnell et al, December 2021 https://www.lacedandlethal.com/

UW Medicine

13

14

Fentanyl- Mexis, Oxy 80s, Percs



- Most street drugs these days will contain fentanyl
- Fentanyl can come in multiple forms and multiple colors
- Samples may look similar but have different ranges of purity and mixes of other substances
- You cannot tell what is in a drug or how strong it is by its appearance

UW Medicine

The Dangers of the Unregulated Drug Market: King County Snapshot 2/15/23



• Powder samples: 15.5 to 30.3% fentanyl





• Pill Samples: Approx 100mg with fentanyl ranging from 1.7%-3.2% (Acetaminophen 27.7%-45.4%)







UW Medicine

15 16

Reasons for Increase in Deaths



- __
- Huge Impact of fentanyl
 - Availability
 - Price
 - Concentration
 - Variability in product
- Perceived lower risk of smoking pills over IVDU
- Covid pandemic isolation
 - Using alone
 - Reduction of regular recovery supports

UW Medicine

Xylazine



- Xylazine is a veterinary anesthetic, not approved for use in humans, often cut into powder heroin/fentanyl, sometimes called trang, trang dope or sleep cut
- It's not an opioid. It has strong sedative effects and excessive sleepiness without euphoria.
- It starts quickly and makes people fall out hard for about the first hour if they're not used to it.
- Xylazine can extend the duration effect of fentanyl, so people don't get sick again in two hours

National Harm Reduction Coalition Harmreduction.org

UW Medicine

17

18

Xylazine Impacts



- Xylazine injection creates vasoconstriction causing hideous <u>necrotic wounds</u> which can become extensive, chronic and difficult to heal on their own
- These can occur beyond the site of injection, e.g. anywhere with a bite or cut Risk of hypo and hyperthermia and injuries from falls due to intense sedation
- Naloxone is not effective on Xylazine but will still help with opioid part of overdose

UW Medicine

During 2019-2020, overdose deaths involving illicitly manufactured fentanyls (IMF) increased sharply in several parts of the country

AMONG IMF-INVOLVED 2020 DEATHS

Wany routes of drug use were reported including injection, snorting, smoking, and ingesting

Dis. Jy/MMW/R7050e3

MMW/R

https://www.cdc.gov/stopoverdose/naloxone/index.html

19

20

What is Naloxone



- Opioid overdose antidote, an antagonist- blocks the opioid receptor to restore breathing
- Has no effect on someone who has not taken opioids and is safe to use on children and pregnant women.
- Effect lasts 30-90 minutes only
- Does not make someone more likely to use more drugs
- It is rare someone reacts aggressively after receiving naloxone

UW Medicine



21

22

Tips for Talking about Naloxone



_

- Discuss other risk reduction strategies
- Help the client make an OD prevention plan
- Make sure others know where they keep naloxone
- Take the opportunity to talk about protecting friends and family members if appropriate

UW Medicine



23





